


Anterior Cruciate Ligament Reconstruction



32

KYLE HOPPE & ANGELA LAUTEN
DPT CANDIDATES 2013

Objectives



- To understand risk factors associated with ACL injury
- To understand normal anatomy & biomechanics of the ACL as it relates to rehab activities
- To gain new insight into the surgical procedure: graft selection & fixation
- To understand healing properties of tissues & how this affects the rehab timeline

Objectives



- To understand the need for criterion-based progression
- Describe functional outcome measures appropriate for patients following ACLR
- List the factors necessary for consideration to start return-to-play training

The Need



- 41% of athletes returning to prior level of activity (Davies, 2013; Arden et al., 2011)
 - Performance reduced by 1/3rd when returning from ACLR
- 4-fold increase in re-injury following ACLR (Creighton et al., 2010)
- Lack of standardization/“best” practice (Creighton et al., 2010)
- 13% of 264 articles report using outcome measures (Barber-Westin & Noyes, 2011)
- 40-90% develop OA within 10 years (Davies, 2013)

The Issues



- 20-25% of patients experience unsatisfactory results at 7-10yrs following ACLR (Jung et al, 2009)
 - 15-25% continue to suffer pain & instability
- ACL disruption alters knee kinematics (Dargel et al, 2007)
- Current ACLR techniques do not seem to fully restore normal physiology of movement (Getgood & Spalding, 2012)
- Multifactorial causes of ACLR failure (Hosseini et al, 2012)
- Rehab protocols are irrespective of age & graft selection (Heijne & Werner, 2007)

Considerations




- Risk factors for ACL injury
- Mechanisms/factors in injury
- Anatomy & biomechanics of the ACL
- Key factors influencing the outcome of ACLR
 - Graft selection
 - Graft fixation
 - Rate of graft healing
 - Optimal post-op rehab?

Risk Factors for ACL Injury (Smith et al., 2012)



- Neuromuscular
 - Different postures & landing biomechanics
 - Reduced core proprioception
- Anatomic
 - Knee geometry
 - ✦ Intercondylar notch width
 - Generalized/knee laxity
 - Static alignment



Risk Factors for ACL Injury (Smith et al., 2012)



- **Intrinsic risk factors**
 - Gender differences
 - Hormones
 - Genetic
 - Previous injury
- **Extrinsic risk factors**
 - Weather
 - Type & condition of playing surface
 - Footwear



Mechanisms & Factors in Injury (Dargel et al., 2007)




- Active quad pull
- Isolated disruption vs concomitant injuries
 - Lesions to medial/lateral meniscus as well as to MCL reported in ~80% of cases
 - Minor/major bruising of chondral or subchondral structures
- Concept of primary & secondary restraints

Anatomy/Physiology of the ACL



- Dense connective tissue containing parallel rows of fibroblasts & type I collagen
- Originates from posteromedial aspect of lateral femoral condyle
- Inserts into anterolateral aspect of medial tibial spine


Knee - Cruciate and Collateral Ligaments
Right Knee in Flexion: Anterior View




F. Netter
M.D.

Anatomy/Physiology of the ACL

- Divided into 2 bundles:
 - Anteromedial (AM) bundle
 - ✦ Mean length = 33mm
 - Posterolateral (PL) bundle
 - ✦ Mean length = 18mm



21



b



13

Biomechanics of the ACL



- **Primary restraint**
 - Anterior translation of tibia on femur
 - Hyperextension
 - Rotation (IR of tibia on femur)
- **Secondary restraint:**
 - Varus/valgus
- **Each bundle has different kinematic roles** (Gabriel et al., 2004)
 - AM bundle = more taut in knee flex (~90deg)
 - PL bundle = more taut in knee ext (<30deg)
 - In-situ forces of each bundle under different loads

Biomechanics of the ACL



Biomechanics of the ACL




- Native femur-ACL-tibia complex (FATC) (Woo et al., 2006)
 - Ultimate failure load avg: 2160 ± 157 N
 - Stiffness avg: 242 ± 28 N/mm
- Effects of aging (Woo et al., 1991)
 - Ultimate failure load avg: 495 ± 85 N
 - Stiffness avg: 124 ± 16 N/mm

Effects of specimen age and orientation on the structural properties of the FATC (mean \pm SEM)

Age group	Specimen orientation	Stiffness (N/mm)	Ultimate load (N)	Energy absorbed (N-m)
Younger (22–35)	Anatomical	242 ± 28	2160 ± 157	11.6 ± 1.7
	Tibial	218 ± 27	1602 ± 167	8.3 ± 2.0
Middle (40–50)	Anatomical	220 ± 24	1503 ± 83	6.1 ± 0.5
	Tibial	192 ± 17	1160 ± 104	4.3 ± 0.5
Older (60–97)	Anatomical	180 ± 25	658 ± 129	1.8 ± 0.5
	Tibial	124 ± 16	495 ± 85	1.4 ± 0.3

Biomechanics of Normal Walking (Shelburne et al., 2005)




- Pattern of force in the ACL explained almost entirely by anterior pull of patellar tendon
- Intact knee
 - Peak ACL force occurs at midstance, ~303N
 - ✦ Force explained by balance of muscle forces, joint contact forces, & ground reaction forces (GRF) applied to the leg
- ACL-deficient knee
 - Increase in anterior tibial translation
 - ✦ Effects on MCL force

Goal of ACLR



- To restore ACL function & normal knee joint kinematics
- Anatomical restoration = functional restoration of the ACL to its native dimensions, collagen orientation, & insertion sites (Yabroudi & Irrgang, 2013)



Key Factors that Influence the Outcome of ACLR



GRAFT SELECTION

GRAFT FIXATION

RATE OF GRAFT HEALING

OPTIMAL POST-OP REHAB?

Graft Selection



Autografts

- Bone-patellar-tendon bone (BPTB)
- Hamstring (HS) grafts:
 - Double stranded
 - Quadruple stranded
- Quadriceps tendon

Allografts

- BPTB
- HS
- Tibialis anterior tendon
- Tibialis posterior tendon
- Achilles tendon
- Fascia lata

Autografts: BPTB Advantages



- High strength & stiffness
- Consistency of size of graft
- Bone plugs on both ends of graft – facilitates better fixation & incorporation
- High success rate
- No risk of disease transmission
- No immunogenic response

Autografts: BPTB Disadvantages



- Harvest site morbidity:
 - Anterior knee pain, pain when kneeling
 - Possible patellar fx or late patellar tendon rupture
- Quad weakness
- Extensor deficit
- Patellofemoral pain
- Loss of motion



Autografts: HS grafts



Advantages

- Less donor site morbidity?
- Less anterior knee pain
- No extensor mechanism dysfunction

Disadvantages

- No pre-op assessment of graft thickness
 - Variability in HS size
- Soft tissue to bone healing (slower rate)
- Knee flexor strength deficits
- Graft tunnel motion or enlargement

Allografts: General Advantages



- No associated donor site morbidity
- No weakening of extensor/flexor apparatus
- Shorter operating time (no graft harvest)
- Larger and predictable graft sizes
- Greater availability
 - Appropriate for revision surgery, multi-ligament injury, or in the presence of patellar baja
- Less pain

Allografts: General Disadvantages



- Delayed graft incorporation
- Immunogenic response of the host to the graft
- Risk of disease transmission (bacterial/viral)
- Sterilization techniques (Baer & Harner, 2007)
 - Radiation & high failure rates
 - Compromising properties of graft – doses above 2Mrad
- Graft cost
- Large age-related differences
 - Structural properties of graft

Graft References



Autografts

- Dheerendra et al, 2012
- Dargel et al, 2007
- Karlsson et al, 2011
- Woo et al, 2006
- Hsu et al, 2010
- Baer & Harner, 2007
- Meuffels et al, 2012

Allografts

- Dheerendra et al, 2012
- Prodromos et al, 2007
- Sun et al, 2009
- Baer & Harner, 2007
- Woo et al, 1991
- Meuffels et al, 2012
- Gulotta & Rodeo, 2007

Biomechanical properties of grafts (Baer & Harner, 2007)



<u>Graft Type</u>	<u>Ultimate Tensile Load</u>	<u>Stiffness</u>	<u>Cross-Sectional Area (CSA)</u>
<i>Native ACL</i>	<i>~2160 N</i>	<i>242 N/mm</i>	<i>44mm²</i>
BPTB auto	2977 N	620 N/mm	35mm ²
HS auto: quadruple strand	4090 N	776 N/mm	53mm ²
Tib Ant allo	4122 N	460 N/mm	48.2mm ²
Tib Post allo	3594 N	370 N/mm	44.4mm ²
Achilles tendon allo	4617 N	685 N/mm	67mm ²

Is one graft better than the other?



- Several studies have demonstrated comparable results between the grafts in terms of laxity, IKDC scores, & return to activity
 - Barrett et al., 2005
 - ✦ BPTB auto vs. allo, pts 40yo & older – both graft choices highly effective
 - Rihn et al., 2006
 - ✦ BPTB auto vs allo, pts 25-40yo – 94% (autos) & 95% (allos) rated their activity levels as normal/nearly normal

Is one graft better than the other? (Mohtadi et. al, 2011)



- 2011 Cochrane review – BPTB vs. HS autografts
 - No difference in re-rupture rate or IKDC scores
 - All tests for knee stability favored BPTB grafts
 - BPTB group: knee ext loss; HS group: knee flex loss
 - Conclusion:
 - ✦ Current evidence insufficient to recommend either of the 2 grafts for ACLR

Indications/Recommendations for Graft Selection (Vyas et al., 2012)



- **Depends on several factors:**
 - Pt preference, age, activity level, physical requirements, expected outcomes, timeline for return to play, associated ligament injuries, medical comorbidities, previous surgery, tissue availability, & surgeon preference/experience
- **Common indications for autograft:**
 - Young, very active pts
 - Pts 30-40yo who are highly involved in athletics
- **Common indications for allograft:**
 - Mild-mod active pt >40yo w/ symptomatic instability
 - Multi-ligament reconstruction or revision ACLR
 - Cases where autograft tissue is inadequate
 - Pt preference

BOTTOM LINE (Baer & Harner, 2007)



- No graft choice can match completely the characteristics & function of the native ACL
- The ideal graft choice should:
 - Have similar biomechanical properties to that of the native ACL
 - Have low morbidity
 - Incorporate quickly
 - Be able to restore functional stability to the knee over the *long term*
 - ✦ Taking into account: pt preference, activity level, prior surgery, comorbidities, & goals!

Key Factors that Influence the Outcome of ACLR



GRAFT SELECTION

GRAFT FIXATION


RATE OF GRAFT HEALING

OPTIMAL POST-OP REHAB?

Graft Fixation (Dargel et al., 2007; Brand et al., 2000)



- Fixation devices generally fail to reconstruct complex nature of native FATC
- A graft & its fixation devices are loaded, therefore displacement is affected by direction of force & stiffness of graft+fixation
 - Bungee-cord effect
 - ✦ Longitudinal
 - Windshield-wiper effect
 - ✦ Transverse
- Fixation = weakest link early on



Key Factors that Influence the Outcome of ACLR



GRAFT SELECTION

GRAFT FIXATION


RATE OF GRAFT HEALING

OPTIMAL POST-OP REHAB?


Healing (Baer & Harner, 2007)



- All grafts undergo a sequential process of healing & ligamentization
 - Inflammation & graft necrosis
 - ✦ 1-2mos after surgery
 - Revascularization & cell repopulation
 - ✦ 20 days to 6mos after surgery
 - ✦ Changes in material properties of graft
 - Remodeling
 - ✦ May take up to 18mos to occur



25



25

Bone-to-bone Healing (Gulotta & Rodeo, 2007)




- Incorporation relatively quick: 6-8wks
 - At 6wks, weakest point becomes the junction of the graft bone plug & native insertion of patellar tendon (if autograft)
- Components of tendon-to-bone healing
- Factors that may impede healing:
 - Length of tendon portion in BPTB grafts is greater than the intra-articular ACL length
 - Presence of synovial fluid at graft-tunnel interface

Tendon-to-bone Healing (Gulotta & Rodeo, 2007)



- Takes considerably longer, often 8-12wks to mature
- Strength of soft-tissue bone interface may increase most during 1st 4wks after ACLR
- Factors that may impede healing:
 - Presence of inflammation at graft site = scar formation
 - Slow/limited bone ingrowth into tendon graft
 - Graft-tunnel motion
 - Insufficient # of undifferentiated progenitor cells at healing interface – age!
 - Lack of coordinated signaling cascade



Allograft Tissue Healing



- **Slower rate than autograft tissue** (Baer & Harner, 2007)
 - Prolonged inflammatory stage, delayed remodeling of collagen fibers
- **Strength of fixations & maturation of graft are delayed** (Escamilla et al., 2012; Gulotta & Rodeo, 2007)
 - Greater decrease in implantation structural properties
 - Slower rate of biologic incorporation
- **Remodeling phase – graft loses tensile strength initially** (Gulotta & Rodeo, 2007)
- ****Accelerated protocols should be utilized with caution** (Escamilla et al., 2012; Vyas et al., 2012)

Factors in Healing & Considerations for Rehab



- **Age-related changes** (Woo et al., 1991)
 - Properties of ligaments/tendons can change with advancing age & activity level
- **Stress/loading**
 - Ligaments & grafts remodel according to applied motion & stress
 - Graft fixation remains the weak link in early post-op period
- **Blood supply**
- **Graft type**

Key Factors that Influence the Outcome of ACLR



GRAFT SELECTION

GRAFT FIXATION

RATE OF GRAFT HEALING

OPTIMAL POST-OP REHAB?

Optimal Rehab? (Yabroudi & Irrgang, 2013)



- **Depends on:**
 - Surgical procedure
 - Concomitant surgical procedures performed
 - Pt's previous level of activity/fitness
 - Response to surgery & rehab
 - Desired activity level after surgery
- **Progression affected by several factors:**
 - Surgical procedure: anatomic/nonanatomic?
 - Graft type
 - Presence of associated injuries
 - Pt variation

Goals of ACLR Rehab (Yabroudi & Irrgang, 2013)



- **Several goals remain constant:**
 - Early ROM
 - Preservation of quad function
 - Progression of functional activities
 - Not exceeding limits of the involved tissue's healing properties

Goals of ACLR Rehab (Yabroudi & Irrgang, 2013)



Table 1
Summary of the primary goals for each stage of rehabilitation after anatomic ACL reconstruction

Early Postoperative	Strengthening and Neuromuscular Control	Return to Activity and Sports
Controlling pain and edema	Progression of strengthening	Complete the entire functional rehabilitation spectrum
Protecting the healing graft	Neuromuscular control	
Minimizing the effects of immobilization	Improving balance	Make a full return to the patient's previous level of daily, occupational and athletic activity, and sport participation
Obtaining full passive and active extension of the knee symmetric to the noninvolved knee	Preparation for the return to activity and sports stage	
Achieving 100° to 120° of knee flexion		
Preserving quadriceps muscle function		
Restore the ability to perform straight-leg raise without a quadriceps lag		
Progression to full weight bearing		
Achieving normal gait		

Optimal Rehab? More things to Consider...

- Effect of quad activity on ACL loading
 - Relative increase
- Influence of knee flex angle
 - The “danger zone”:
0-30deg
- Hamstring co-contraction
 - Not effective at small knee flex angles



General Progression



- **Early Rehabilitation Goals** (Myer et al., 2006)
 - ROM
 - Decrease swelling/effusion
 - Patellar Mobility (Wilk et al., 2012)
 - Gait
 - Body Weight Strengthening Exercises
 - ✦ Quad set, SLR, Heel slides
 - Strength Training: OKC vs CKC

*Controlled, with ROM & strengthening guidelines

General Progression



- **Late Rehabilitation Activities** (Myer et al., 2006)
 - Jogging/Running
 - Jumping/Hopping
 - Sport Specific Drills (cutting, shuffling)

*Less restricted, leading to variation in practice

Accelerated vs. Nonaccelerated



- **Early weight-bearing and ROM** (Beynnon et al, 2011; DeCarlo et al, 1992)
 - Laxity the same or better in Accelerated
- **Brace in extension** (Melegati et al, 2003)
- **Addition of sport-specific drills** (Wright et al, 2008; Beynnon et al, 2011; DeCarlo et al, 1992; Roi et al, 2005)
- **Criterion-based progression** (Roi et al, 2005)

OKC versus CKC



- **Minimize strain on the healing graft** (Yu and Garrett, 2007):
 - 1st 3 months: Limited arc of motion between 90-60 flexion for OKC; 0-60 flexion for CKC
 - 3-4 months post: OKC thru full arc as tolerated by the joint
 - OKC flexion can increase hamstring strength
 - ✦ Avoid in 1st 4-6 weeks if meniscus is also repaired
- **Eclectic Approach combining both OKC and CKC**
(Glass et al, 2010)

Why Criterion-Based Progression?

(Davies, 2013; Roi et al, 2005; Myer et al, 2006)



- Provides objective measures that allow for patients to see deficits compared to subjective comments/feelings
- Justifies rationale for sport restriction
- Evidence for payers
- Sets benchmarks/goals for athletes to strive for



Progression Criteria



- Factors that *should* influence decision making:
 - Sex
 - Age
 - PMHx
 - Prior surgeries
 - Current surgery (graft selection, technique)
 - SxS
 - Motivation
 - Psychological health of athlete
 - Imaging studies
 - Concomitant injuries

Progression Criteria



- **Pain/Swelling: incomplete healing if present** (Creighton et al, 2010)
- **ROM:**
 - 0-100 degrees in initial 2 weeks (Wilk et al, 2012)
 - Full ROM before intense strength training/sports drills
- **Strength Training: No extensor lag** (Myer et al, 2006; Wilk et al, 2012)
- **Normal Gait** (Myer et al, 2006)
- **Dynamic Stability** (Wilk et al, 2012; Davies, 2013; Myer et al, 2006)

Initiation of Sport Specific Drills

(Davies, 2013)



- Physician's okay
- Patient's subjective comments/pain ratings
- Stage of tissue healing/strength of graft
- Patient's objective scores WNL
- Rehabilitation path

Initiation of Sport Specific Drills



- Negative Pivot Shift/No patient report of "giving away" (Myer et al, 2006)
- IKDC >70 (Myer et al, 2006)
- Quad Peak Torque/Body Weight (@180 deg/sec):
60-65% males; 50-55% females (Wilk et al, 1994)
- Dynamic Stability during ADLs/ambulation (Myer et al, 2006; Wilk et al, 2012)



What to Work on



- **Single limb strength and stability** (Fitzgerald et al, 2000)
 - Single limb squats or lunges
 - Surface Modifications (Wilk et al, 2012)
- **Treadmill running/walking** (Myer et al, 2006)
 - Improve symmetry
- **Squats** (Myer et al, 2006; Ohkoshi et al, 1991)
 - Even distribution (<20% difference)
- **Landing on single limb**
- **Endurance**
- **Plyometric Training** (Myer GD, Ford KR et al, 2006)

Outcome Measures



- ROM
- Swelling
- Isokinetic Strength

Functional Tests:

- Y-Balance Test
- Star Excursion Balance Test
- Hop Tests

Patient Reported:

- IKDC
- Cincinnati Knee Rating Scale
- KOOS

Return-to-Play:


Tegner

Psychometric Properties of Outcome Measures for ACLR



- **International Knee Documentation Committee (IKDC):** 3-subscales measuring subjective symptoms, sport/activity, and function (Irrgang et al, 2009)
- **Cincinnati Knee Rating System:** 13-section patient report of pain, swelling, sports, and work (Barber-Westin et al, 1999)
- **Knee Injury and Osteoarthritis Outcome Score (KOOS):** 5-subscales that measure: Pain, Symptoms, ADLs, Sport/Rec, and QoL (Salavanti et al, 2011; Wang et al, 2010)
- **Tegner Activity Scale:** 11-point scale to rate level of activity/sport participation (Briggs et al, 2009)

Star Excursion Balance Test



Y-Balance Test




Figure 1. The three reach directions of the Lower Quarter Y Balance Test (a. anterior, b. posteromedial, c. posterolateral).

Hop Tests




- 85-90% of uninjured side (Davies, 2013; Bise, 2013)


Single Hop for Distance




6-m Timed Hop



Triple Hop for Distance



Crossover Hop for Distance



Return-to-Play

(Davies, 2013; Creighton et al, 2010)



- Considerations:
 - Type of Sport
 - ✦ High, Moderate, or Low Risk
- Clinician Risk
- External Pressures



Return-to-Play Criteria



- **The “best” we have:**
 - ROM (Creighton et al, 2010; Barber-Westin and Noyes, 2011)
 - ✦ 70-100%, Full ROM, Pain-free
- **Laxity** (Davies, 2013; Barber-Westin and Noyes, 2011; DeCarlo, 1992)
 - 3mm (MAX) 2.5 or less preferred on Lachman or KT measurement
- **Thigh Circumference** (Creighton et al, 2010; Barber-Westin and Noyes, 2011)
 - <1cm difference
 - No Swelling or Effusion
 - Measurement sites

Return-to-Play Criteria




- **Isokinetic Strength Measures** (Barber-Westin and Noyes, 2011; Wilk et al 1994)
 - <10-15% side-to-side difference
- **Functional Stability** (Barber-Westin and Noyes, 2011; Davies 2013; Reid et al, 2007)
 - Functional Test Scores
 - Single-Leg Hop Tests (>85%-90%)
- **Psychological Confidence** (Creighton, 2010; Chmielewski, 2008)
 - Fear most common reason for not returning to sport

CONCLUSIONS



- Individual characteristics
- Biomechanics
- Graft types
- Healing of tissue properties
- Individual treatment and rehab
 - Outcome measures
 - Criterion-based progression



THANK YOU!!



ANY QUESTIONS, PLEASE CONTACT:

ALAUTEN@MED.UNC.EDU

KYLE_HOPPES@MED.UNC.EDU

References



1. Andersson D, Samuelsson K, Karlsson J. Treatment of anterior cruciate ligament injuries with special reference to surgical technique and rehabilitation: an assessment of randomized controlled trials. *Arthroscopy*. 2009;25(6):653-85.
2. Ardern CL, Webster KE, Taylor NF, Feller JA. Return to sport following anterior cruciate ligament reconstruction surgery: a systematic review and meta-analysis of the state of play. *Br J Sports Med*. 2011 Jun;45(7):596-606.
3. Baer GS, Harner CD. Clinical Outcomes of Allograft Versus Autograft in Anterior Cruciate Ligament Reconstruction. *Clin Sports Med*. 2007;26(4):661-81.
4. Barber-Westin SD, Noyes FR, McCloskey JW. Rigorous statistical reliability, validity, and responsiveness testing of the Cincinnati knee rating system in 350 subjects with uninjured, injured, or anterior cruciate ligament-reconstructed knees. *Am J Sports Med*. 1999 Jul-Aug;27(4):402-16.
5. Barber-Westin, SD, Noyes, FR. Factors used to determine return to unrestricted sports activities after ACL-R. *Arthroscopy*. 27:1697-1705, 2011
6. Barrett G, Stokes D, White M. Anterior cruciate ligament reconstruction in patients older than 40 years: allograft versus autograft patellar tendon. *Am J Sports Med* 2005;33(10):1505-12.
7. Beynon, B. D., Johnson, R. J., Naud, S., Fleming, B. C., Abate, J. A., Brattbakk, B., et al. (2011). Accelerated versus nonaccelerated rehabilitation after anterior cruciate ligament reconstruction: A prospective, randomized, double-blind investigation evaluating knee joint laxity using roentgen stereophotogrammetric analysis. *The American Journal of Sports Medicine*, 39(12), 2536-2548
8. Brand J, Weiler A, Caborn D, Brown C, Johnson D. Graft Fixation in Cruciate Ligament Reconstruction. *Am J Sports Med*. 2000;28(5):761-74.
9. Briggs KK, Lysholm J, Tegner Y, Rodkey WG, Kocher MS, Steadman JR. The reliability, validity, and responsiveness of the Lysholm score and Tegner activity scale for anterior cruciate ligament injuries of the knee: 25 years later. *Am J Sports Med*. 2009 May;37(5):890-7.
10. Chmielewski, LT, Jones D, Day, T, Tillman, SM, Lentz TA, George SZ. The association of pain and fear of movement/reinjury with function during anterior cruciate ligament reconstruction rehabilitation. *J Orthop Sports Phys Ther*. 2008;38:746-753.
11. Cohen SB, Sekiya JK. Allograft safety in anterior cruciate ligament reconstruction. *Clin Sports Med*. 2007;26(4):597-605.
12. Creighton, DW, et.al. Return-to-play in sport: a decision-based model. *Clin J Sports Med*. 20(5):379-385, 2010
13. Dargel J, Gotter M, Mader K, Pennig D, Koebke J, Schmidt-Wiethoff R. Biomechanics of the anterior cruciate ligament and implications for surgical reconstruction. *Strategies Trauma Limb Reconstr*. 2007;2(1):1-12.
14. Davies, G. Functional testing algorithm and clinical decision making for return to play criteria for the lower extremity (Presentation). Combined Sections meeting. 2013; San Diego, California.
15. De Carlo, M., Shelbourne, K. D., McCarroll, J. R., & Rettig, A. C. (1992). Traditional versus accelerated rehabilitation following ACL reconstruction: A one-year follow-up. *Journal of Orthopaedic & Sports Physical Therapy*, 15(6), 309-316.

References



16. DeMorat G, Weinhold P, Blackburn T, Chudik S, Garrett W. Aggressive Quadriceps Loading Can Induce Noncontact Anterior Cruciate Ligament Injury. *Am J Sports Med.* 2004;32(2):477-83.
17. Dheerendra SK, Khan WS, Singhal R, Shivarathre DG, Pydisetty R, Johnstone D. Anterior cruciate ligament graft choices: a review of current concepts. *Open Orthop J.* 2012;6:281-6.
18. Escamilla RF, Macleod TD, Wilk KE, Paulos L, Andrews JR. Anterior cruciate ligament strain and tensile forces for weight-bearing and non-weight-bearing exercises: a guide to exercise selection. *J Orthop Sports Phys Ther.* 2012;42(3):208-20.
19. Fitzgerald GK, Axe MJ, Snyder-Mackler L. The efficacy of perturbation training in nonoperative anterior cruciate ligament rehabilitation programs for physical active individuals. *Phys Ther.* 2000;80:128-140.
20. Gabriel MT, Wong EK, Woo SL, Yagi M, Debski RE. Distribution of in situ forces in the anterior cruciate ligament in response to rotatory loads. *J Orthop Res.* 2004;22(1):85-9.
21. Getgood A, Spalding T. The evolution of anatomic anterior cruciate ligament reconstruction. *Open Orthop J.* 2012;6:287-94.
22. Ghodadra NS, Mall NA, Grumet R, et al. Interval arthrometric comparison of anterior cruciate ligament reconstruction using bone-patellar tendon-bone autograft versus allograft: do grafts attenuate within the first year postoperatively?. *Am J Sports Med.* 2012;40(6):1347-54.
23. Glass R, Waddell J, Hoogenboom B. The Effects of Open versus Closed Kinetic Chain Exercises on Patients with ACL Deficient or Reconstructed Knees: A Systematic Review. *N Am J Sports Phys Ther.* 2010;5(2):74-84.
24. Grodski M, Marks R. Exercises following anterior cruciate ligament reconstructive surgery: biomechanical considerations and efficacy of current approaches. *Res Sports Med.* 2008;16(2):75-96.
25. Gulotta LV, Rodeo SA. Biology of autograft and allograft healing in anterior cruciate ligament reconstruction. *Clin Sports Med.* 2007;26(4):509-24.
26. Harner CD, Olson E, Irrgang JJ, et al. Allograft versus autograft anterior cruciate ligament reconstruction: 3- to 5-year outcome. *Clin Orthop Relat Res* 1996;324:134-44.
27. Heijne A, Werner S. Early versus late start of open kinetic chain quadriceps exercises after ACL reconstruction with patellar tendon or hamstring grafts: a prospective randomized outcome study. *Knee Surg Sports Traumatol Arthrosc.* 2007;15(4):402-14.
28. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med.* 2005;33(4):492-501.
29. Hosseini A, Lodhia P, Van de velde SK, et al. Tunnel position and graft orientation in failed anterior cruciate ligament reconstruction: a clinical and imaging analysis. *Int Orthop.* 2012;36(4):845-52.
30. Hsu SL, Liang R, Woo SL. Functional tissue engineering of ligament healing. *Sports Med Arthrosc Rehabil Ther Technol.* 2010;2:12.
31. <http://blog.brianschiff.com/?tag=hip-strength>

References



32. <http://blog.nasm.org/tag/acl-reconstruction/>
33. http://jill-adayinthelife.blogspot.com/2008_11_01_archive.html
34. <http://moon.ouhsc.edu/dthompso/namics/pasfor.htm>
35. <http://thetimes-tribune.com/news/health-science/acl-rehab-tough-part-begins-1.1293692>
36. <http://www.msdlatinamerica.com/ebooks/PracticalOrthopaedicSportsMedicineArthroscopy/sid538403.html>
37. [http://www.orthopaedicsone.com/display/Main/Revision+Anterior+Cruciate+Ligament+\(ACL\)+Reconstruction](http://www.orthopaedicsone.com/display/Main/Revision+Anterior+Cruciate+Ligament+(ACL)+Reconstruction)
38. <http://www.texarkanagazette.com/news/2013/01/08/rookie-mistake-rg3-staying-in-game-may-r-686744.php>
39. Irrgang JJ, Anderson AF, Boland AL, et al. Development and validation of the international knee documentation committee subjective knee form. *Am J Sports Med.* 2001;29:600-613.
40. Jung HJ, Fisher MB, Woo SL. Role of biomechanics in the understanding of normal, injured, and healing ligaments and tendons. *Sports Med Arthrosc Rehabil Ther Technol.* 2009;1(1):9.
41. Kaeding CC, Aros B, Pedroza A, et al. Allograft Versus Autograft Anterior Cruciate Ligament Reconstruction: Predictors of Failure From a MOON Prospective Longitudinal Cohort. *Sports Health.* 2011;3(1):73-81.
42. Karlsson J, Irrgang JJ, Van eck CF, Samuelsson K, Mejia HA, Fu FH. Anatomic single- and double-bundle anterior cruciate ligament reconstruction, part 2: clinical application of surgical technique. *Am J Sports Med.* 2011;39(9):2016-26.
43. Leys T, Salmon L, Waller A, Linklater J, Pinczewski L. Clinical results and risk factors for reinjury 15 years after anterior cruciate ligament reconstruction: a prospective study of hamstring and patellar tendon grafts. *Am J Sports Med.* 2012;40(3):595-605.
44. Li G, Defrate LE, Rubash HE, Gill TJ. In vivo kinematics of the ACL during weight-bearing knee flexion. *J Orthop Res.* 2005;23(2):340-4.
45. Li H, Tao H, Cho S, Chen S, Yao Z, Chen S. Difference in graft maturity of the reconstructed anterior cruciate ligament 2 years postoperatively: a comparison between autografts and allografts in young men using clinical and 3.0-T magnetic resonance imaging evaluation. *Am J Sports Med.* 2012;40(7):1519-26.
46. Markatos K, Kaseta MK, Lалlos SN, Korres DS, Efstathopoulos N. The anatomy of the ACL and its importance in ACL reconstruction. *Eur J Orthop Surg Traumatol.* 2012;[Epub ahead of print].
47. Melegati, G., Tornese, D., Bandi, M., Volpi, P., Schonhuber, H., & Denti, M. (2003). The role of the rehabilitation brace in restoring knee extension after anterior cruciate ligament reconstruction: A prospective controlled study. *Knee Surgery, Sports Traumatology, Arthroscopy: Official Journal of the ESSKA*, 11(5), 322-326. doi:10.1007/s00167-003-0386-3.
48. Meuffels DE, Poldervaart MT, Diercks RL, et al. Guideline on anterior cruciate ligament injury. *Acta Orthop.* 2012;83(4):379-86.
49. Mohtadi NG, Chan DS, Dainty KN, Whelan DB. Patellar tendon versus hamstring tendon autograft for anterior cruciate ligament rupture in adults. *Cochrane Database Syst Rev.* 2011;(9):CD005960.
50. Myer GD, Ford KR, Brent JL, Hewett TE. The effects of plyometric vs. dynamic stabilization and balance training on power, balance, and landing force in female athletes. *J Strength Cond Res.* 2006 May;20(2):345-53.
51. Myer GD, et al. Rehabilitation after ACL Reconstruction: criteria-based progression through return-to-sport phase. *JOSPT.* 36:403-414, 2006

References



52. Ohkoshi Y, Yasuda K, Kaneda K, Wada T, Yamanaka M. Biomechanical analysis of rehabilitation in the standing position. *Am J Sports Med.* 1991;19:605-611.
53. Perez, F, Vicen, J. Effects of Power Balance® in the static and dynamic balance in physically active subjects. *Apunts. Medicina de l'Esport.* 2011;46(171): 109-115.
54. Pflum MA, Shelburne KB, Torry MR, Decker MJ, Pandy MG. Model prediction of anterior cruciate ligament force during drop-landings. *Med Sci Sports Exerc.* 2004;36(11):1949-58.
55. Prodromos C, Joyce B, Shi K. A meta-analysis of stability of autografts compared to allografts after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2007;15(7):851-6.
56. Reid A, Birmingham TB, Stratford PW, Alcock GK, Giffin JR. Hop testing provides a reliable and valid outcome measure during rehabilitation after anterior cruciate ligament reconstruction. *Phys Ther.* 2007 Mar;87(3):337-49.
57. Rihn JA, Irrgang JJ, Chhabra A, et al. Does irradiation affect the clinical outcome of patellar tendon allograft ACL reconstruction? *Knee Surg Sports Traumatol Arthrosc* 2006;14(9): 885-96.
58. Roi, G. S., Creta, D., Nanni, G., Marcacci, M., Zaffagnini, S., & Snyder-Mackler, L. (2005). Return to official italian first division soccer games within 90 days after anterior cruciate ligament reconstruction: A case report... including commentary by shelbourne KD and fithian DC with author response. *Journal of Orthopaedic & Sports Physical Therapy*, 35(2), 52-66.
59. Salavati M, Akhbari B, Mohammadi F, Mazaheri M, Khorrami M. Knee injury and Osteoarthritis Outcome Score (KOOS); reliability and validity in competitive athletes after anterior cruciate ligament reconstruction. *Osteoarthritis Cartilage.* 2011 Apr;19(4):406-10.
60. Shelburne KB, Pandy MG, Anderson FC, Torry MR. Pattern of anterior cruciate ligament force in normal walking. *J Biomech.* 2004;37(6):797-805.
61. Shelburne KB, Torry MR, Pandy MG. Muscle, ligament, and joint-contact forces at the knee during walking. *Med Sci Sports Exerc.* 2005;37(11):1948-56.
62. Siegel L, Vandenakker-Albanese C, Siegel D. Anterior cruciate ligament injuries: anatomy, physiology, biomechanics, and management. *Clin J Sport Med.* 2012;22(4):349-55.
63. Smith HC, Vacek P, Johnson RJ, et al. Risk factors for anterior cruciate ligament injury: a review of the literature - part 1: neuromuscular and anatomic risk. *Sports Health.* 2012;4(1):69-78.
64. Smith HC, Vacek P, Johnson RJ, et al. Risk factors for anterior cruciate ligament injury: a review of the literature-part 2: hormonal, genetic, cognitive function, previous injury, and extrinsic risk factors. *Sports Health.* 2012;4(2):155-61.
65. Sun K, Tian S, Zhang J, Xia C, Zhang C, Yu T. Anterior cruciate ligament reconstruction with BPTB autograft, irradiated versus non-irradiated allograft: a prospective randomized clinical study. *Knee Surg Sports Traumatol Arthrosc.* 2009;17(5):464-74.
66. Vyas D, Rabuck SJ, Harner CD. Allograft anterior cruciate ligament reconstruction: indications, techniques, and outcomes. *J Orthop Sports Phys Ther.* 2012;42(3):196-207.

References



67. Wang D, Jones MH, Khair MM, Miniaci A. Patient-reported outcome measures for the knee. *J Knee Surg.* 2010 Sep; 23(3):137-51.
68. Wilk KE, Marcina LC, Cain EL, et al: Recent advances in the rehabilitation of ACL injuries. *J Orthop Sports Phys Ther.* 2012;42:208-220.
69. Wilk KE, Romaniello WT, Soscia SM, Arrigo CA, Andrews JR. The relationship between subjective knee scores, isokinetic testing, and functional testing in the ACL-reconstructed knee. *J Orthop Sports Phys Ther.* 1994;20:60-73.
70. Woo SL, Hollis JM, Adams DJ, Lyon RM, Takai S. Tensile properties of the human femur-anterior cruciate ligament-tibia complex. The effects of specimen age and orientation. *Am J Sports Med.* 1991;19(3):217-25.
71. Woo SL, Wu C, Dede O, Vercillo F, Noorani S. Biomechanics and anterior cruciate ligament reconstruction. *J Orthop Surg Res.* 2006;1:1-9.
72. Wright, R. W., Preston, E., Fleming, B. C., Amendola, A., Andrish, J. T., Bergfeld, J. A., et al. (2008). A systematic review of anterior cruciate ligament reconstruction rehabilitation: Part II: Open versus closed kinetic chain exercises, neuromuscular electrical stimulation, accelerated rehabilitation, and miscellaneous topics. *The Journal of Knee Surgery*, 21(3), 225-234.
73. Yabroudi MA, Irrgang JJ. Rehabilitation and return to play after anatomic anterior cruciate ligament reconstruction. *Clin Sports Med.* 2013;32(1):165-75.
74. Yasuda K, Van eck CF, Hoshino Y, Fu FH, Tashman S. Anatomic single- and double-bundle anterior cruciate ligament reconstruction, part 1: Basic science. *Am J Sports Med.* 2011;39(8):1789-99.
75. Yu B, Garrett WE. Mechanisms of non-contact ACL injuries. *Br J Sports Med.* 2007;41 Suppl 1:i47-51.