

Sources for Patient Handouts¹⁻¹³

1. Skelly AC, Chou R, Dettori JR, et al. *Noninvasive Nonpharmacological Treatment for Chronic Pain: A Systematic Review*. Rockville (MD): Agency for Healthcare Research and Quality (US); 2018. doi:10.23970/AHRQEPCCER209.
2. Allen K, Choong P, Davis A, et al. Osteoarthritis: Models for appropriate care across the disease continuum. *Best Pract Res Clin Rheumatol* 2016;(30):503-535.
3. Fransen M, McConnell S, Harmer AR, Van der Esch M, Simic M, Bennell KL. Exercise for osteoarthritis of the knee: a Cochrane systematic review. *Br. J. Sports Med.* 2015;49(24):1554-1557. doi:10.1136/bjsports-2015-095424.
4. Shakoor N, Sengupta M, Foucher KC, Wimmer MA, Fogg LF, Block JA. Effects of common footwear on joint loading in osteoarthritis of the knee. *Arthritis Care Res. (Hoboken)* 2010;62(7):917-923. doi:10.1002/acr.20165.
5. Kerrigan DC, Johansson JL, Bryant MG, Boxer JA, Della Croce U, Riley PO. Moderate-heeled shoes and knee joint torques relevant to the development and progression of knee osteoarthritis. *Arch. Phys. Med. Rehabil.* 2005;86(5):871-875. doi:10.1016/j.apmr.2004.09.018.
6. Beumer L, Wong J, Warden SJ, Kemp JL, Foster P, Crossley KM. Effects of exercise and manual therapy on pain associated with hip osteoarthritis: a systematic review and meta-analysis. *Br. J. Sports Med.* 2016;50(8):458-463. doi:10.1136/bjsports-2015-095255.
7. Fernandes L, Hagen KB, Bijlsma JWJ, et al. EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis. *Ann. Rheum. Dis.* 2013;72(7):1125-1135. doi:10.1136/annrheumdis-2012-202745.
8. American Academy of Orthopaedic Surgeons. MANAGEMENT OF OSTEOARTHRITIS OF THE HIP EVIDENCE-BASED CLINICAL PRACTICE GUIDELINE. 2017.
9. American Academy of Orthopaedic Surgeons. TREATMENT OF OSTEOARTHRITIS OF THE KNEE: EVIDENCE-BASED GUIDELINE, 2ND EDITION. 2013.
10. HEP2go - Build a HEP < Home Exercise Program> For Free. Available at: https://www.hep2go.com/index_b.php?userRef=gciaake. Accessed March 1, 2020.
11. Vieira S, Dibai-Filho AV, Brandino HE, Ferreira VTK, Scheicher ME. Abdominal muscle strength is related to the quality of life among older adults with lumbar osteoarthritis. *J Bodyw Mov Ther* 2015;19(2):273-277. doi:10.1016/j.jbmt.2014.05.002.
12. Chen H, Onishi K. Effect of home exercise program performance in patients with osteoarthritis of the knee or the spine on the visual analog scale after discharge from physical therapy. *Int. J. Rehabil. Res.* 2012;35(3):275-277. doi:10.1097/MRR.0b013e328355a1bd.
13. Adams MA, Lama P, Zehra U, Dolan P. Why do some intervertebral discs degenerate, when others (in the same spine) do not? *Clin Anat* 2015;28(2):195-204. doi:10.1002/ca.22404.

*Exercise images borrowed from hep2go.com